

Primitive Shoshonites from Fiji: Mineralogy, Melt Inclusions and Geochemistry

Roman A.J. Leslie



UNIVERSITY
OF TASMANIA

Submitted in fulfilment of the requirements for the degree of
Doctor of Philosophy

December 2004

Declaration

This thesis contains no material which has been accepted for the award of any other degree or diploma in any tertiary institution and to the best of my knowledge and belief, contains no material previously published or written by another person, except where due reference is made in text.



R.A.J. Leslie

3rd Dec. 2004

Access Authority

This thesis may be made available for loan and limited copying in accordance with the copyright Act, 1968.


R.A.J. Leslie
3rd Dec. 2004.

Abstract

Fundamental issues regarding the origin and evolution of primitive shoshonitic magmas are addressed using mineralogical, melt inclusion and geochemical data from Fijian shoshonites. Melt inclusions in olivine phenocrysts from primitive Fijian shoshonites are used to critically assess the issues relating to melt sampling by high-Fo olivine phenocrysts and address to what degree trapped melt compositions reflect the larger volume magmatic system as a whole.

Shoshonitic magmas erupted in Fiji during the Pliocene (5-3Ma) from 11 main volcanic centres along three broad ENE and NNW trending lineaments. The most mafic shoshonitic lavas (absarokites) range from 8.4-15.2wt% MgO and are highly porphyritic, containing phenocrysts of olivine (to Fo_{93.2}) and clinopyroxene (to Mg# 93.3).

The vast majority of melt inclusions in high-Fo-olivine phenocrysts from Fijian shoshonites have anomalous major element composition, mainly characterised by high CaO contents and high-CaO/Al₂O₃. Anomalous melt inclusion compositions are interpreted to reflect localised, grain-scale dissolution-reaction-mixing processes within the magmatic plumbing system where hot, primitive magma comes in contact with wall-rocks and/or pre-existing semi-solidified mush zones. Injection of hot primitive melt causes partial dissolution of the mush-zone phases, which are not in equilibrium with the primitive melt and mixing of the reaction products with the primitive magma. Rapid cooling at the margins of the magma body induces fast crystallisation and efficient trapping of numerous and large melt inclusions, with anomalous major element composition. Populations of melt inclusions in high-Fo olivine phenocrysts from Fijian shoshonites, and arguably many other subduction-related suites, are naturally biased toward anomalous compositions.

The rare occurrence of normal melt inclusions in high-Fo olivine allows reconstruction of primitive shoshonitic melt compositions. Estimated parental

shoshonitic melts in equilibrium with olivine Fo_{92.6} have ~16wt% MgO and evolve to compositions ~3-4wt% MgO, via ~63% total crystallisation, involving olivine, Cr-spinel, clinopyroxene, magnetite, plagioclase with minor apatite, phlogopite and/or K-feldspar.

High-Fo olivine phenocrysts (>Fo₈₅) crystallised from primitive shoshonitic melt batches, then resided in cumulate layers within the magmatic system for several months, as inferred from modelling of Fe-loss diffusion profiles surrounding melt inclusions. High-Fo olivine phenocrysts were subsequently entrained from cumulate layers by variably evolved shoshonitic melts during the eruption process. Absarokites with primitive shoshonitic compositions (high MgO contents) consist of variably evolved transporting melts and a cargo of crystals, often derived from different parts of the magmatic system. In most cases the crystal cargo bears no direct relationship to the liquid(s) that entrain and transport them

Fijian shoshonitic suites display a range of enrichment in LILE, Th and P relative to REE and HFSE, reflecting variable contributions by subduction-related components. These components are: a sediment-melt component that, following dehydration of subducted sediment, transfers K, Th, P₂O₅, LREE, some Sr and U to the sub-arc mantle; fluid(s) derived from the dehydration of both subducted oceanic crust and sediment that transfers predominantly Pb, Sr, Ba, K and U; and the sub-arc mantle wedge source component that controls HFSE, HREE, to a lesser degree LREE.

Fijian shoshonites are derived partial melting of the metasomatised, non-convecting upper mantle. Melt production, concentrated at sites of maximum metasomatism in the non-convecting sub-Fijian mantle, focused strain and caused nucleation of lithosphere-scale transtensional or strike-slip shear zones, facilitating rapid rotation of the Fiji Platform between 5-3Ma. Primary shoshonitic melts represent hybrid, vein + wall rock melts, which segregated from upper mantle at ~1.5GPa (~50km depth) with an average melt fraction of ~7% and a temperature of ~1330°C. Advective heating and partial melting of metasomatised sub-arc mantle beneath Fiji occurred in response to a vigorous in-flow of hot asthenosphere following detachment of the stalled Pacific plate at ~12Ma along the Vitiaz trench.

Acknowledgements

This project would not have been possible without the ongoing support and encouragement of Leonid Danyushevsky, who has gone far beyond the role of supervisor, to make this challenging experience into a rewarding one. Thanks to my co-supervisor Tony Crawford for keeping me well constrained and giving me focus. I'd also like to thank the Minerals and Resources Department in Fiji for permission to undertake field work in Fiji, Noby, at Ovalau Watersports, Dere Bay Resort, Koro and to the people of the Islands of Koro, Moala, Kadavu for giving me permission to take samples from their land.

At the University of Tasmania, I'd particularly like to thank Phil Robinson, Katie McGoldrick, David Steele, Dima Kamenetsky, June Pongratz, Darren Andrews, Andrew Wurst and Kieren Howard, Neil Martin; although innumerable others have helped in some way. A special thanks goes to the morning coffee crew who've kept me sane.

Thankyou so much to my Mum and Dad, Julia and Robert, for your patience and belief in me. Finally, thankyou Kate and Oliver for providing support and some comic relief during this often not comic process.

Contents

Declaration

Access Authority

Abstract

Acknowledgements

Abbreviations

Chapter 1 Introduction.....	1
1.1 Aims.....	1
1.2 Structure.....	3
Chapter 2 Shoshonitic magmas.....	4
2.1 The shoshonite magma series: background and nomenclature.....	4
2.2 Generation of potassic magmas.....	6
2.3 Tectonic setting of the shoshonitic rock association.....	13
2.3.1 Arc settings.....	14
2.3.2 Post-collision settings.....	17
2.3.3 Obscure/ambiguous settings.....	18
2.4 Summary.....	19
Chapter 3 Melt inclusion studies related to the study of primitive magmas.....	20
3.1 Melt trapping in the magmatic system.....	21
3.2 The study of melt inclusions.....	25
3.2.1 Post-entrapment modification.....	25
3.2.2 Reconstructing trapped melt compositions-“the ideal case”.....	26
3.2.2.1 Reheating melt inclusions.....	26
3.2.2.2 Glassy inclusions.....	28
3.2.2.3 Modelling of olivine crystallisation within melt inclusions.....	29
3.2.3 Reconstructing trapped melt compositions-“the non-ideal case”.....	30
3.2.3.1 Hydrogen diffusion and H ₂ O dissociation.....	30
3.2.3.2 Fe-loss in olivine hosted melt inclusions.....	33
3.2.3.3 Modelling of crystallisation and re-equilibrium in olivine-hosted melt inclusions.....	34
Chapter 4 Tectonic setting and geological evolution of Fiji.....	41
4.1 Regional setting.....	41
4.2 Plate tectonic history.....	45
4.3 Crustal structure and present-day tectonics.....	49
4.3.1. The sub-Fijian mantle: a graveyard for subducted slabs.....	51
4.4 The geological evolution of Fiji.....	52

4.4.1 Early arc stage (>10 Ma).....	52
4.4.2 Mature arc stage (10-5 Ma).....	55
4.4.3 Early rifting stage (5-3 Ma).....	57
4.4.4 Late rifting stage (~3.0 Ma-present).....	63
4.5 Summary.....	68
Chapter 5 Shoshonitic magmatism in Fiji.....	70
5.1.1 The geology of Tavua Volcano.....	70
5.1.2 The geology of the Astrolabe Islands.....	76
5.1.3 The geology of Vatu-i-Cake.....	82
5.1.4 The geology of Moala.....	85
5.2 Geochemistry of Fijian shoshonitic suites.....	86
5.2.1 Geochemical classification.....	86
5.2.2 Major elements.....	86
5.2.3 Trace elements.....	97
5.2.4 Isotopic compositions.....	99
5.3 Summary.....	102
Chapter 6 Mineralogy.....	103
6.1 Sample selection and preparation.....	103
6.2 Olivine.....	104
6.2.1 Introduction and petrographic characteristics.....	104
6.2.2 Olivine major element composition and population distribution.....	109
6.2.3 Major element (Fe-Mg) zoning in olivine phenocrysts.....	110
6.2.4 Minor and trace elements in olivine.....	111
6.2.4.1 Minor elements in olivine from Tavua Volcano.....	111
6.2.4.1.1 Minor element zoning in Tavua olivine.....	115
6.2.4.2 Minor elements in olivine from the Astrolabe Group.....	116
6.2.4.3 Minor elements in olivine from Vatu-i-Cake.....	119
6.3 Spinel inclusions in olivine.....	121
6.3.1 Introduction.....	121
6.3.2 Spinel inclusions in olivine from Tavua Volcano.....	121
6.3.2.1 Classification and subdivision of spinel-series minerals.....	122
6.3.2.2 Tavua spinel compositions.....	122
6.3.2.3 Coexisting spinel inclusions.....	128
6.3.2.4 Tavua spinel summary.....	129
6.3.3 Spinel inclusions in olivine from the Astrolabe Group and Vatu-i-Cake.....	129
6.3.3.1.1 Astrolabe Group spinel compositions.....	129
6.3.3.1.2 Astrolabe Group chromite compositions.....	133
6.3.3.2 Vatu-i-Cake spinel compositions.....	134

6.3.3.3 Summary of Astrolabe Group and Vatu-i-Cake spinel compositions.....	136
6.4 Clinopyroxene.....	
6.4.1 Clinopyroxene phenocrysts.....	136
6.4.1.1 Composition and population distribution.....	137
6.4.1.2 Compositional variation.....	143
6.4.2 Olivine-hosted clinopyroxene inclusions.....	144
6.4.3 Enstatite inclusions in Vatu-i-Cake olivine.....	145
6.5 Plagioclase inclusions in Tavua olivine.....	145
6.6 Groundmass compositions.....	145
6.7 Summary of mineralogy.....	147
Chapter 7 Melt inclusions.....	149
7.1 Petrography.....	149
7.1.1 Tavua.....	149
7.1.2 Astrolabe Group.....	154
7.1.3 Vatu-i-Cake.....	155
7.1.4 Fluid halos and fluid loss in olivine-hosted melt inclusions.....	156
7.2 Experimental techniques and results.....	161
7.2.1 Sample preparation.....	161
7.2.2 Heating stage.....	161
7.2.3 Experimental procedure and summary.....	165
7.2.4 Assessment of experimental results.....	168
7.3.1 Fe-loss in olivine-hosted melt inclusions.....	177
7.3.2 Fe-loss diffusion profiles.....	177
7.3.2.1 Melt inclusion EL9/OL-7/137m1.....	179
7.3.2.2 Melt inclusion EL9OL-4/269.....	188
7.3.2.3 Melt inclusion EL9/OL-4/263.....	189
7.3.2.4 Melt inclusion EL9/OL-5/9m2.....	190
Chapter 8 Anomalous melt inclusion compositions.....	192
8.1 The composition of melt inclusions from Fijian shoshonites.....	192
8.2 Inclusion compositions hosted by magnesian olivine (>Fo83).....	196
8.2.1. Anomalous inclusion compositions from Fijian shoshintes.....	197
8.3 Other examples of anomalous melt inclusion compositions and explanations to account for their origin.....	200
8.3.1 Comparison with other subduction related suites.....	202
8.3.2 Examples of MORB: ophiolite geology and anomalous melt inclusion compositions in phenocrysts from MORB.....	205

8.4 Magmatic plumbing, crystallisation conditions and dissolution-reaction-mixing processes.....	208
8.5 Primitive melt-wall rock/mush zone interaction and the origin of anomalous inclusion compositions in Fijian shoshonites.....	211
8.5.1 DRM processes and the origin of High-CaO inclusion compositions.....	211
8.5.2 DRM processes and the origin of low-CaO inclusion compositions.....	215
8.5.3 DRM processes and crystalline (solid) inclusions in high-Fo olivine.....	216
8.5.4 DRM processes and minor element concentrations in olivine.....	222
8.6 The trace element geochemistry of anomalous inclusion compositions.....	226
8.6.1 The geochemical effects of clinopyroxene and apatite assimilation.....	228
8.7 Discussion.....	233
Chapter 9 Primitive shoshonitic melts from Fiji.....	240
9.1 The major element composition of primitive Fijian shoshonitic melts.....	240
9.1.1 Correction for Fe-loss and $\text{Fe}^{2+}/\text{Fe}^{3+}$ determination.....	240
9.1.2 Major elements and liquid line of descent.....	247
9.1.3 Estimating a parental shoshonitic melt composition.....	249
9.1.4 Degree of fractionation.....	252
9.1.5 Summary and discussion.....	255
9.2 Volatiles.....	257
9.2.1 Chlorine.....	257
9.2.2 Fluorine.....	259
9.2.3 Sulphur.....	261
9.2.3.1 Sulphur speciation.....	264
9.2.4 H_2O concentrations.....	266
9.2.5 Summary of volatile data.....	268
9.3 Trace element geochemistry.....	270
9.3.1 Normal inclusion compositions from Tavua.....	270
9.3.2 Geochemical variations between Fijian shoshonitic suites.....	272
9.3.3 Chalcophile elements (Cu and Zn).....	276
9.4 Trace element and volatile abundance of the estimated parental melt composition.....	281
Chapter 10 The origin and evolution of shoshonitic magmas in Fiji.....	283
10.1 The origin of shoshonitic magmas in Fiji.....	284
10.1.1 Introduction.....	284
10.1.2 The composition of the unmodified (pre-subduction) Fijian mantle.....	286
10.1.3 Fijian shoshonitic magmas: an expression of subduction-related magmatism.....	290
10.1.4 Inter-suite variations among Fijian shoshonites and the nature of the subduction-derived components.....	293

10.1.4.1 There are shoshonites and there are shoshonites: distinguishing between and subduction-derived components (fluids and melts) in Fijian shoshonites.....	302
10.1.5 Generating shoshonitic melts.....	304
10.1.6 Relationship to regional geodynamics and tectonics.....	308
10.1.6.1 Thermal constraints.....	311
10.1.6.2 Tectono-magmatic evolution.....	314
10.2 Cooling history and rock evolution.....	316
Chapter 11 Conclusions.....	330
References	